

A service of the National Library of Medicine
and the National Institutes of HealthMy NCBI
[Sign In] [Register]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

for **Breast cancer recurrence menstrual cycle**

[Go] [Clear]

Limits

Preview/Index

History

Clipboard

Details

About Entrez
NCBI Toolbar

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-UtilitiesPubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBIRelated Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central**Note:** Performing your original search, **Breast cancer recurrence menstrual cycle**, in PubMed will retrieve [63 citations](#).

Display Citation

Show 20

Sort by

Send to

All: 1

Review: 0

**1:** [Ann Intern Med.](#) 1991 Sep 1;115(5):337-42.

Related Articles, Links

Comment in:

- [Ann Intern Med. 1991 Sep 1;115\(5\):401-3.](#)

Timing of breast cancer excision during the menstrual cycle influences duration of disease-free survival.**Senie RT, Rosen PP, Rhodes P, Lesser ML.**

Centers for Disease Control, Atlanta, Georgia.

OBJECTIVE: To study disease-free survival at 10 years in relation to timing of breast tumor excision during the menstrual cycle. **DESIGN:** A prospective study of consecutively treated patients with primary breast cancer. **SETTING:** Memorial Sloan-Kettering Cancer Center, New York. **PATIENTS:** Two hundred and eighty-three premenopausal patients treated by mastectomy and axillary dissection. **MAIN RESULTS:** When the tumor was excised during the follicular phase, approximated by setting the putative day of ovulation on day 14 after the onset of last menses, a higher recurrence risk (43%) was observed compared with excision later in the menstrual cycle (29%, $P = 0.02$). The rate peaked among patients treated between days 7 and 14 and was lowest between days 20 and 30. Multivariate analysis using the Cox regression model to control for tumor size, nodal status, estrogen receptor status, adjuvant chemotherapy, and family history indicated that the hazard rate of breast cancer recurrence after excision during the follicular phase was 1.53 (95% CI, 1.02 to 2.29). Stratification by nodal status indicated that the effect of phase was statistically significant only among patients with positive nodes (hazard ratio, 2.10; CI, 1.19 to 3.70). **CONCLUSIONS:** Our results support the hypothesis that the risk for recurrence may be affected by the hormonal milieu of the menstrual cycle; these findings must be confirmed, however, by a prospective study in which cycle phase at time of tumor excision is biochemically documented.

MeSH Terms:

- [Adult](#)
- [Breast Neoplasms/surgery*](#)
- [Carcinoma/secondary](#)
- [Carcinoma/surgery](#)
- [Female](#)
- [Humans](#)
- [Lymphatic Metastasis](#)
- [Mastectomy, Modified Radical/methods*](#)
- [Menstrual Cycle/physiology*](#)
- [Middle Aged](#)
- [Proportional Hazards Models](#)
- [Prospective Studies](#)
- [Statistics](#)
- [Survival Rate](#)
- [Time Factors](#)

PMID: 1863022 [PubMed - indexed for MEDLINE]

Display [Citation](#)

Show 20 Sort by

Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Apr 24 2006 06:33:44